

16. Rodent Management in Thailand

Puangtong Boonsong, Sermsakdi Hongnark, Kornkaew Suasa-ard, Yuvaluk Khoprasert, Prasarttong Promkerd, Greangsak Hamarit, Piyanee Nookarn and Thomas Jäkel

Abstract

Thailand is an agricultural country where rice and other agricultural products contribute a substantial part to the gross domestic product. Rodents cause problems both in agriculture and as reservoirs of human diseases. Some basic data on the agricultural situation in Thailand and the damage inflicted by rodents as well as an overview of past and present efforts in rodent control are presented. The Ministry of Agriculture and Cooperatives is responsible for rodent problems in agriculture. In the Department of Agriculture, the Agricultural Zoology Research Group conducts research on rodent problems. It devises methods for rodent control and is responsible for the transfer of technical know-how to extension programs. Some aspects of the group's research are highlighted, including the use of the endemic parasitic protozoan *Sarcocystis singaporensis* as a potential biocontrol agent. As the Thai government aims to substantially reduce the use of pesticides, integrated pest management concepts, including new strategies in rodent control, are being pursued in pilot areas.

Keywords

Thailand, rodent management, rodent research, integrated pest management, rodent borne diseases, biological control, *Sarcocystis singaporensis*, ecology

INTRODUCTION

THAILAND IS A tropical country located in Southeast Asia surrounded by Cambodia, Lao People's Democratic Republic, Myanmar and Malaysia. The peninsula runs down to the Indian Ocean, and receives the south-west monsoon from mid-May to October. It covers a land area of 513,178 square kilometres and extends about 1,620 kilometres from north to south and 775 kilometres from east to west. There are three seasons: cool from November to February, summer from March to May and rainy season from June–October. The average minimum temperature is 20°C and average maximum temperature is 37°C. Annual rainfall averages from 1,000–2,000 mm, varying greatly from place to place and year to year.

Thailand is divided into four regions. The *northern region* is mostly mountain highlands where many rivers originate and run down to the central plain. In the north, agriculture is mostly limited to the fertile valleys of the Chao Phraya River tributaries. Fruit trees, forest trees and vegetable crops are the main sources of income in the region. The *north-eastern region* has a flat rolling terrain called the Khorat Plateau. Much of the land has poor soil fertility and little water. Large areas are flooded during the rainy season but are very dry during the rest of the year. Provinces along the Mae Kong River use this water for agriculture with rice being cultivated mainly for home consumption. Fruit trees and rubber are being promoted to help green the area. The *central plain* is regarded as the rice bowl of Thailand. Corn,

fruit crops and vegetables are also of economic significance in the region. The *southern region* has several sizeable coastal plains and a mountain chain running along its western coast. This region has mostly sandy loam soil suitable for fruit trees and tree crops, especially rubber and oil palm.

MAIN CROPS, AREAS INVOLVED AND CONTRIBUTION TO THE ECONOMY

Thailand cultivates about 26,523,836 ha or 51.7% of its total land area (Anonymous 1996). Irrigated land comprises about 15.3% of the agricultural area. Of this area, rice comprises about 52%, field crops 25%, fruit trees and tree crops 16% and idle land 2.5%. The remaining land is grassland, housing and other areas. Rice has a farm value of nearly US\$3 billion. Other major field crops are cassava, corn, sugarcane, oil crops, perennial trees such as rubber, and fruit trees (Table 1).

The importance of agriculture to the Thai economy can be measured by its contribution to the gross domestic product (GDP). Agriculture comprises 16% of the GDP, industry 24%, and commerce and service sectors the remaining 60%. Apart from its contribution to the GDP, agriculture boosts the national economy through wealth distribution and provides gainful employment for approximately 64% of the Thai population.

CROP DAMAGE BY RODENTS

Major pest species

Although there are about 33 murid species in the southeastern end of the Asian continent (Corbet and Hill 1992), less than half of these

are considered pests in Thailand. The two main ecotypes of rodents found in temperate zones, those occurring in grassland and woodland (Wood 1994), also largely apply to the situation in Thailand. There are pests of field crops and those of forestry and orchards. Additionally, cosmopolitan species like *Rattus norvegicus* are also prevalent. Table 2 lists the key pest species of various crops as observed by the Agricultural Zoology Research Group (AZRG) of the Department of Agriculture during field surveys (Ratanaworabhan 1971; Suasa-ard et al. 1987; Khoprasert et al. 1990; Hongnark et al. 1994).

Damage in lowland

Rodent problems in lowland occur mostly in rice and field crops in the central regions of Thailand. Because the Chao Phaya River and Tha Chin River run through the area and

70% of the region is irrigated, farmers can cultivate throughout the year. In Suphan Buri, Nakhon Pathom and Pathum Thani, rice varieties are cultivated which allow five harvests every two years; alternatively, field crops (soybean, mungbean, baby corn etc.) are grown after harvesting the major rice crop. When food is available all year, rodents can breed throughout the whole period (Boonsong et al. 1984a).

The importance of the problem of damage to rice by rodents in Thailand previously led to the introduction of a nationwide control scheme by the Thai-German Rodent Control Project (see below). At that time (1976–77), damage assessment in rice was performed in central, southern, northern, and north-eastern Thailand according to established methods (Weis 1981).

Table 1.
Area planted and estimated farm value of principal crops in 1995/96 in Thailand (rate of exchange 1 US\$ = 40 baht).

Crop	Area (million ha)	Farm value (US\$ million)
Rice	0.14	2622.1
Rubber	1.82	1064.0
Fruits	0.60	659.1
Vegetables	0.30	527.1
Coconut	0.38	78.0
Oil palm	0.10	115.6
Soy bean	0.30	83.5
Sugar cane	1.00	559.4
Maize	1.33	420.7
Cassava	1.26	426.0
Mungbean	0.35	69.5

Source:
Agricultural Statistics of Thailand, Crop year 1995/1996. Office of Agricultural Economics, Ministry of Agriculture and Cooperatives

In each of the four regions, three provinces were randomly selected and three districts in each province inspected. In each district, percentage damage was measured on eight plots (30 m × 30 m each) two weeks before harvest of the wet-season rice. Figure 1 shows that, on average, about 18% of the rice was damaged in the central plains which translates to losses of approximately US\$300 million. A more recent survey (1990–93) by the AZRG, employing the same methods in the same areas, showed that the situation in rice had improved (Figure 1) (Hongnark et al. 1993) although an average of 1.5% damage still equates to losses of about US\$35 million. Whether this reduction of the problem in rice can be entirely attributed to the control scheme (see below) or is in part due to other factors such as natural fluctuations in rodent populations is not known. Certainly, awareness of control measures among farmers has increased substantially. Currently,

problems with rodents in rice appear to be moderate. It should be noted that *Rattus argentiventer* which is considered the most serious rice field pest besides *Bandicota indica* in Thailand (Wood 1994) was observed rarely during recent surveys of AZRG. Instead, *Rattus losea* seemed to be more abundant.

In oil palm plantations, losses caused by rats vary considerably both between years and between plantations. Damage to mature palms generally ranged from 6–36% (Boonsong et al. 1987). Rodents infesting older plantations are climbing species which prefer ripe oil palm fruits. Younger oil palms are attacked by ground-dwelling species (Table 2). Although conspicuous damage was patchy, trapping showed that rats are well spread (Wood 1987). The density of *Rattus tiomanicus* was reported to range from about 125–625 rats/ha in Malaysia (Wood and Liau 1984), and the situation appears to be similar in Thailand (Wood 1987).

Table 2.
Major rodent pest species in Thailand and the crops/areas they affect.

Species	Rice fields and field crops	Oil palms	Fruit trees (Mango, longan, macadamia etc.)	Storage and houses
<i>Bandicota indica</i>	✓	✓ (young palms, < 3 years)		
<i>Bandicota savilei</i>	✓			
<i>Rattus argentiventer</i>	✓	✓ (young palms, < 3 years)		
<i>Rattus bowersi</i>		✓ (young palms, < 3 years)		
<i>Rattus exulans</i>				✓
<i>Rattus losea</i>	✓		✓	
<i>Rattus norvegicus</i>				✓
<i>Rattus rattus</i>	✓	✓	✓	✓
<i>Rattus tiomanicus</i>		✓		
<i>Mus caroli</i>	✓		✓	
<i>Mus cervicolor</i>	✓		✓	
<i>Mus musculus</i>				✓




Figure 1. Average pre-harvest damage (%) in rice in southern (S), northern (N), north-eastern (NE), and central (C) Thailand, and the average (AV) of the four regions from 1990–1993 compared with 1976–1977 (Hongnark et al. 1993).

Damage in upland

Upland is defined as those areas that are 600 m above sea level. In Thailand, most upland is in the north, with some in the north-eastern and southern regions. In these areas, rice, wheat, barley, temperate fruits (apple, pear, strawberry, macadamia etc.), vegetables (cabbage, carrot, onion, broccoli etc.) and coffee are cultivated. In upland, rice damaged by rodents appears to be generally low. Birds such as spotted munia (*Lonchura punctulata*), sharp-tailed munia (*Lonchura striata*) and Pegu sparrow (*Passer flaveolus*) are more problematic (Hongnark et al. 1984). In the north-east, four species of rodents

(*B. indica*, *R. losea*, *Mus cervicolor*, *Mus caroli*) are a problem in wheat (damage 6.4%, Hongnark et al. 1994). In the north, these species—and additionally *Rattus rattus*—occur in barley plantations which mainly serve the brewing industry. Various plots in a field of 800 ha were damaged by 0.4–17% during harvest (Artchawakom et al. 1986).

RODENTS AS CARRIERS OF DISEASE

A recent outbreak of leptospirosis in a rural area in north-eastern Thailand killed 107 people between October–December 1997 while a total of 2,236 had to be treated for leptospirosis during that year (Chokvivat 1998). The incident was broadly covered in the media and it drew fresh attention to the rodent problem. Although it is not clear which rodent species actually transmitted the disease, this incident emphasises the need for a better knowledge of the epidemiology of rodent-borne diseases in this region. During 1986–1988, only 466 cases of leptospirosis were reported. This record is believed to underestimate the real incidence of the disease because laboratory facilities were not appropriate for screening of large numbers of blood samples from all parts of Thailand. Frequently, leptospirosis is incorrectly diagnosed as influenza or a virus infection (Silapapochakul 1992). As a consequence of the recent outbreak of leptospirosis, the Environmental Health Bureau (Department of Health) started an extension program (three years, 1998–2000) to monitor and control leptospirosis in north-eastern Thailand. The activities include (i) training of officials of the Department of Health working in regional health centres in appropriate protection

against leptospirosis, (ii) training of local technicians to work on rodent population estimation in urban areas, and (iii) training in correct and efficient use of rodenticides. Also, universities in Thailand (i.e. Mahidol) started to focus on diseases transmitted by rodents.

B. indica in Thailand and *R. norvegicus* were found to be infected with species of the hantavirus family (Schmaljohn and Hjelle 1997), members of which are responsible for an increasing global health problem. The Seoul virus carried by Norway rats is known to induce haemorrhagic fever with renal syndrome in humans, a condition which can be potentially fatal (Schmaljohn and Hjelle 1997). Sera collected from wild rodents (*B. indica*, *M. cervicolor*, *R. losea* and *R. rattus*) caught in rural areas of Chiang Rai Province, northern Thailand, were positive using an immunofluorescent antibody test (Lietmeyer 1988).

These two examples of rodent-borne diseases are especially relevant to Thailand and form only a part of the long list of pathogens in rodents that can affect humans. This subject has been dealt with in detail previously (Gratz 1994; Singleton and Petch 1994; Schrag and Wiener 1995). It must be noted that humans in Thailand do not only have close contact with rodents (be it in the rice field or inside houses), but additionally, rodents form a part of the diet of the rural population (see also below). Of 142 and 123 farmers who were surveyed for their rodent control practice in two different regions in northern Thailand, 79% and 55%, respectively, reported consuming rodents regularly (Boonsong et al. 1994). In particular, *B. indica* is highly regarded as a meat source.

CURRENT CONTROL METHODS AND EXTENT OF ADOPTION BY FARMERS

Since the Thai–German Rodent Control Project was set up in 1975, systematic, preventative rodent control methods have been recommended for use in rice and field crops (Weis 1981). After the project ceased in 1979, the Department of Agricultural Extension continued promoting the methods in all regions by establishing the Rodent Control Campaign Project (1988–1993). The objective of the project was to decrease rodent populations in agricultural areas and to maintain low populations by training 2,500 extension officers at the subdistrict level to have a good knowledge of rodent control which they could transfer to 250,000 farmers. The operating area included rice fields in 40 provinces, field crops in 12 provinces, and oil palm and cacao in 12 provinces. The total area was 0.864 million hectares with extension plots (where rodenticides were provided through governmental funds) of 16,000 hectares in 40 provinces. After the first year, the Division of Project and Programme Evaluation, Office of Agricultural Economics, undertook an evaluation of practical adoption by farmers (Anonymous 1989). It revealed that most farmers accepted systematic and preventative rodent control. The survey evaluation showed that all farmers in extension plots were highly accepting of mechanical or physical control techniques, but 33% did not accept use of chronic rodenticides. In extension plots, 67% of farmers accepted use of both acute and chronic rodenticides. Acceptance of these poisons in service areas (those areas that had to buy rodenticides on the private market)

was less than 10%. After this initial evaluation was completed, the Department of Agricultural Extension continued the Rodent Control Campaign Project for five years until 1993.

In rice and field crops, the control method consists of two steps (Weis 1981); knock down of the rat population and subsequent maintenance at low density.

Knockdown step

Chemical control with zinc phosphide or trapping, digging, blanketing or drives. Blanketing or drives are conducted by groups of people who circle an area (about 0.24 ha), cut the vegetation and herd the rats into a small area (2–4 square metres) before they are caught or clubbed.

Maintenance of population at low density

Mechanical or physical methods as mentioned above and chemical control using chronic rodenticides (anticoagulants) such as coumatetralyl, brodifacoum, flocoumafen etc.

After the Rodent Control Campaign Project had ended in 1993, further campaigns were organised from 1995 to 1997 in an area of 864,000 ha near the Kong River in the north-eastern region to control rodent invasions from Lao People's Democratic Republic. Governmental service continued also for every province in that free rodenticides were provided if extension officers had spotted a rodent problem. The AZRG further monitored the control practices of farmers to obtain a realistic view of the degree of adoption of the publicised methods. They interviewed farmers growing soybeans (after the rice crop was harvested) in the north, north-eastern and central

regions. This revealed that about 90% of farmers used zinc phosphide to control rodents when damage was conspicuous (Boonsong et al. 1994, 1995, 1996). Farmers used germinated rice (paddy soaked in water for three nights) or broken rice as bait. The proportion of zinc phosphide mixed with the bait was generally 2–3 times higher than the recommended dose indicating an overuse of acute poison. When farmers observed bait-shyness of rats, they switched to a more attractive bait type such as fresh fish, field crabs (*Somanniathelphusa* spp.) or golden apple snails (*Pomacea canaliculata*) caught in rice fields. In the dry season, when crabs and snails are not abundant, they used mechanical control techniques such as shooting, digging or trapping. As mentioned earlier, many farmers consume rats; in such areas mechanical control over chemical methods were preferred. Farmers also learned that during the booting stage to harvest of rice, most rodents do not take poisoned bait due to the presence of the more attractive rice crop. At that time, farmers usually dug out rat holes instead of applying poison. The surveys further showed that farmers growing rice and field crops did not like to use anticoagulant rodenticides for various reasons. First, the price of anticoagulants was higher than that of zinc phosphide. Second, anticoagulants were sold only in big towns and were not as readily available as zinc phosphide, and third, the effect of chronic anticoagulants was considered too slow.

In conclusion, in the long term, the rodent control scheme has been only partially adopted by farmers in rice fields and field crops. There is a reliance on the use of acute rodenticides despite their limitations

(Prakash 1988), with the only alternative being traditional, mechanical control. Control is usually only considered when the problem is obvious. Preventative measures during periods when rodent numbers are low are the exception rather than the rule, and this situation prevails today.

Rats cause extensive damage to oil palm estates in southern Thailand. In large oil palm estates (>30 hectares) the farm managers generally follow Malaysian plant protection technology from the Palm Oil Research Institution of Malaysia (PORIM). PORIM recommends that control with anticoagulant rodenticides should commence when 5% of the oil palm fruits show fresh damage, and control should be repeated over large areas every six months. During field visits by AZRG to oil palm plantations, it was observed that second-generation anticoagulants like flocoumafen occasionally led to extensive secondary poisoning of predators like barn owls (*Tyto alba*) (AZRG, unpublished observation). In small holdings, most farmers are not interested in rodent control. When the price of fresh fruit is low (less than 2 baht per kilogram), oil palms grow in natural conditions without the use of fertiliser or rodent pest control.

Other control approaches like habitat manipulation and protection of known predators of rodents are regularly proposed to farmers by the Department of Agricultural Extension during extension activities. For instance, farmers employ the former by regularly clearing excessive vegetation on dykes. Measures such as reduction of dyke size are also considered when new fields are designed. Individual farmers reported that they especially take

care of predators like birds of prey or snakes, however, this does not appear to be a common view. Thus to date, an integrated rodent control approach, though pursued by individuals, is not occurring on a broad scale.

RODENT RESEARCH BY THE AZRG

Rodent research in agriculture is primarily the responsibility of the Department of Agriculture, especially the Agricultural Zoology Research Group (AZRG) of the Division of Entomology and Zoology. Research activities focus on various species, such as rodents, bats, birds, crabs, various snails and slugs which are injurious to plants. During the last 20 years, research has been conducted on the following topics:

- ▶ Species identification and density estimation of rodents in economic crops such as rice, maize, soybean, mungbean, oil palm and longan.
- ▶ Life history of key pest species.
- ▶ Ecology: seasonal variations in rodent density in economic crops.
- ▶ Crop damage and loss assessment in rice, oil palm, soybean, maize etc.
- ▶ Chemical control: efficacy of rodenticides in the laboratory and in the field.
- ▶ Integrated pest management: combined application of rodenticides, mechanical control and cultural practices.

Following are some examples of research on the population ecology of pest rodents in Thailand.

The home range length (or maximum diameter of the home range) of rodents in

rice fields has been studied at the Rice Research Station of Pathum Thani using eosin stain (Khoprasert et al. 1977) and mark–release captures in Prachin Buri Province (Somsook et al. 1983). It was found that the maximum radius moved within a week was 90 m and 100 m for *B. indica* and *Bandicota savilei*, respectively. *R. argentiventer* moved a maximum radius of 50 m and *R. losea* 46 m.

The long-distance movement of rodents in rice fields has been studied in an area located in Bang Plama District, Suphan Buri Province (about 100 km north-west of Bangkok) (Boonsong et al. 1984b). Two crops of rice per year were grown with the major crop planted in July and harvested in October and the second crop planted in February and harvested in May. A total of 1,253 rodents were caught in monthly field trips during January to October 1984. They were ear-tagged and released in an area of 8 km². The catch consisted of four species, *R. argentiventer* (51.3%), *R. losea* (18%), *B. indica* (20.4%) and *B. savilei* (10.3%). Only six marked rats were recaptured in the release area; three *B. indica*, two *B. savilei* and one *R. losea*. This low number was in part explained by the extreme trap-shyness of tagged rice-field rats; additionally, farmers conducted intensive rodent control campaigns during the study period. Two great bandicoot rats (*B. indica*) were re-trapped 63 and 95 days after release. They had moved about 1 km from the area where rice had already been harvested to the area where rice would be harvested in the next 2 weeks. Two lesser bandicoot rats (*B. savilei*) were recaptured after 70 days when they had moved about 2 km from a rice crop at tillering stage to a harvesting area. The

single recaptured *R. losea* had moved about 1 km within 40 days from rice at the seedling stage to a harvesting area. In conclusion, it appeared that rodents moved towards areas where rice was being harvested.

In the Central plain, Somsook et al. (1983) studied the population dynamics of the lesser rice-field rat, *R. losea*, from March 1982 to March 1984 in rice fields in Prachin Buri Province (about 300 km east of Bangkok). In this area, floating rice is grown once a year. Rice is planted in June and harvested in December. The rice stubble is left in the field until the following February. During the rainy season, the study area was flooded up to 1.5–2 m from August to October 1983. In 1984, the water level in the rainy season was lower with a maximum of 0.50 m. Rodents were trapped for four nights of each month (800 trap nights). When the water level was high, live-traps were placed on polystyrene sheets. Trapping revealed that *B. indica*, *R. argentiventer* and *R. losea* were present with *R. losea* being the dominant species (90%). The population of *R. losea* showed a clear cyclic pattern with numbers increasing towards the harvest period of rice (Figure 2). Quality and availability of food are certainly major factors influencing the breeding of rodents (Singleton and Petch 1994). It appears that the water level in the rice fields also influenced the population as the numbers of rats were considerably higher in the second wet season (Figure 2) when the water level was lower. Interestingly, *R. argentiventer* could not be trapped in the area when the water was high (1982), but was present during a low water level (1983) indicating that the rats had probably moved out of the area during flooding. Breeding of *R. losea* commenced in September, and most

pregnant females were caught in November and December.

The population dynamics of mice in corn fields were studied in the northern region in Tak Fa District, Nakhon Sawan (240 km north of Bangkok) from May 1986 to September 1991 (Boonsong et al. 1991). In this area, corn was the principal crop which suffered extensive damage from rodents. Mark–release trapping was used to study

the movements and to estimate population density of rodents. A total of 2,200 rodents were caught on 37 occasions. Most of the rodents were mice (96.6%) with 71.8% *M. cervicolor* and 24.8% *M. caroli*. Figure 3 shows the population dynamics of *M. cervicolor*. Similar to the situation in rats, fluctuations in mouse populations reflected the changes in agricultural and climatic conditions. During the dry season (November–May)





Figure 2. Population dynamics of the lesser rice-field rat, *Rattus losea*, in floating rice in the central plains of Thailand from March 1982 to March 1984 (Somsook et al. 1983), showing the numbers of rats trapped in 800 trap nights per month and the monthly average rainfall.

populations increased to 40–90 mice/ha, while in the rainy season (May–October) only 12–19 mice/ha were found. Pregnant females were recorded in the dry as well as in the wet season (Figure 3) indicating that *M. cervicolor* reproduced most of the year except February to May. The survival period of this species in the field was about five months. A home range of 300–400 m² was recorded for *M. cervicolor*.

Some recommendations for control campaigns in Thailand have been derived

from the population studies outlined above. Usually, knockdown of rodent populations starts in the dry season after harvest when less food is available and rats readily accept rodenticide bait containing zinc phosphide. This is also the time when drives and blanketing are conducted. Once the population has decreased in the wet season, chronic poisons and mechanical control are used until the booting stage of the various crops.



BIOLOGICAL CONTROL OF RODENTS USING *SARCOCYSTIS SINGAPORENSIS*

Since 1993, the AZRG and the Department of Parasitology, Hohenheim University, Germany, have cooperated in the framework of a GTZ (Deutsche Gesellschaft für Technische Zusammenarbeit—German Technical Cooperation) project to develop a biological method for rodent control using the apicomplexan protozoan *Sarcocystis singaporensis* which naturally occurs in rats in Southeast Asia. This section provides some background information on the biology of this parasite and considers its possible application against rodent pest species.

Biology and host range of *Sarcocystis singaporensis*

S. singaporensis was discovered by Zaman and Colley (1975) at a time when the obligate two-host life cycle of the sarcosporidia had been recognised (Rommel and Heydorn 1972) and research on this group of parasites was intensive. The original material (sporocysts) was obtained from faeces of a reticulated python (*Python reticulatus*) sold at a butcher's shop in Singapore (Zaman and Colley 1975). Shortly after discovery, Zaman (1976) investigated the intermediate host range of this species and found the parasite to be highly host-specific. Only laboratory rats (*R. norvegicus*) were susceptible to infection through the oral route using sporocysts containing sporozoites, the stage infective for the intermediate host. Other animals, like mice (*Mus musculus*), dogs, cats, chickens, and a rhesus monkey were not susceptible to infection and showed no clinical signs of disease (Zaman 1976; Beaver and Malekar 1981). Infection of rats by

sporozoites is usually followed by two rounds of asexual multiplication inside endothelial cells of various organs, a process by which merozoites are formed (Brehm and Frank 1980). About one month after infection, merozoites eventually invade the muscles to form characteristic cysts (so-called 'sarcocysts') in the striated muscles which contain a third stage, the bradyzoite. Bradyzoites are infective for pythons once the snake preys on rodents.

Subsequently, the definitive and intermediate host range of *S. singaporensis* was studied in more detail using numerous snake and rodent species from various parts of the world (Häfner and Frank 1984; Häfner 1987; Jäkel et al. 1996, 1997b). These studies confirmed the reticulated python in Southeast Asia as the natural definitive host and most suitable to infection with respect to the quantity and quality of sporocysts that developed in the snake's intestine. It appears that *S. singaporensis* also occurs in Australia. Morphologically similar sarcocysts were found in *Rattus fuscipes* (Rzępczyk and Scholtyssek 1976). *Aspidites melanocephalus*, the Australian black-headed python, was found experimentally to be a suitable definitive host (Häfner 1987). The natural definitive host in Australia still remains to be determined. Among boid snakes outside the Australasian region, only *Python sebae*, the African rock python, could be experimentally infected. However, numbers of sporocysts shed with faeces were low and morphological anomalies of sporocysts occurred indicating that these snakes do not provide optimum conditions for the parasite's development (Häfner 1987). Therefore, it seems unlikely that *S. singaporensis* can survive elsewhere, even

if other python species and rats were present.

Among rodents, *Rattus* spp. and *Bandicota* spp. were suitable intermediate hosts (Häfner and Frank 1984). Additionally, *Nesokia indica*, the short-tailed bandicoot rat, was highly susceptible to infection (Jäkel et al. 1996).

Pathogenic effects of *S. singaporensis* in rodents

Zaman (1976) was the first to recognise the pathogenic potential of *S. singaporensis*. He observed that infection of laboratory rats resulted in acute disease, and death, beyond a particular inoculation dose. Wood (1985) made similar observations on infected Malayan wood rats (*R. tiomanicus*) in the laboratory. This was important because it indicated that there existed a parasite with a potential to control wild rats. Up-to-date data on the pathogenicity of parasites in wild rodents are scarce.

The stage responsible for disease in rodents is the merozoite which develops inside endothelial cells. Subclinical infections (which probably prevail in the

wild) are characterised by two distinct peaks of merozoite development in the rat—one occurs around day 6 post infection (p.i.), the other around day 16 p.i. (Brehm and Frank 1980). After inoculation of a lethal quantity of sporocysts, numbers of merozoites increase enormously around day 11 p.i., especially in the lungs. This induces a fatal pneumonia (Jäkel et al. 1996). The factors responsible for the pathology are not fully understood. Mechanical destruction of endothelial cells due to massive development of merozoites seems one likely cause. Furthermore, it has been demonstrated that tumour necrosis factor released by macrophages upon encounter with parasite-antigen is able to kill cultivated cells (Fayer et al. 1988).

The project has determined the degree of pathogenicity of *S. singaporensis* in wild Norway rats (*R. norvegicus*) from Southeast Asia (Thailand), North Africa (Egypt) and Europe (Germany). The parasite appears to be more virulent in hosts occurring outside its natural distribution range. Rats outside Southeast Asia can be killed with about one tenth of the inoculation dose (Table 3).

Table 3. Dose-dependent mortality of wild Norway rats (*Rattus norvegicus*) of different geographic origin after infection with *Sarcocystis singaporensis*^a (original data).

Origin of rats	Inoculation dose (Number of sporocysts)				
	1×10^4	2×10^4	5×10^4	1×10^5	2×10^5
Thailand	n.d. ^b	0/10 (0) ^c	2 (1)/10 (2)	7 (0)/10 (1)	10 (7)/10 (7)
Egypt	0/6	10/10	4/4	n.d.	n.d.
Germany	0/6	14/8	3/3	n.d.	n.d.

^a a parasite isolate from Thailand was used

^b n.d. = not determined

^c numbers of rats that died within 16 days/numbers of rats inoculated. Numbers in parenthesis indicate those rats which were naturally infected.

R. norvegicus is the most resistant species in Thailand. Other species like *Rattus exulans*, *R. argentiventer* and *R. tiomanicus* become moribund at much lower sporocyst doses. Bandicoot rats (*Bandicota* spp.) appear to be particularly susceptible to infection as indicated by massive development of sarcocysts. Adult bandicoot rats usually do not survive an inoculation with 8×10^4 to 1×10^5 sporocysts (AZRG, unpublished observation).

Whether the bradyzoite (the chronic stage inside muscles of the rat) can cause considerable pathologic effects is equivocal. Intriguingly, the number of parasites which develop in striated muscles can be extremely high (up to a billion per gram muscle), especially in wild rats (Figure 4), often without causing any apparent signs of disease. On the other hand, we have observed considerable numbers of laboratory rats as well as wild Norway rats (*R. norvegicus*) that become anorexic, and show rough fur and slow movements during chronic infection. It has been demonstrated that *Sarcocystis* infection renders rodents prone to predation (Hoogenboom and Dijkstra 1987). Impaired mobility due to high sarcocyst numbers in muscle tissue may be an important contributing factor.

There are controversial observations concerning the impact of *S. singaporensis* infection on rodent fecundity. During experiments performed in the last 10 years at the Department of Parasitology, Hohenheim University, Germany, it was observed that sub-lethally infected female laboratory rats (Wistar and F-344 strains) either did not become pregnant or aborted litters. However, in a recent experiment with laboratory-bred wild Norway rats

(*R. norvegicus*) from Germany, infection had no effect on fertility as the number of progeny of infected females was similar to non-infected controls (T. Jäkel, unpublished observation).




Figure 4. Density of bradyzoites of *Sarcocystis singaporensis* in the muscles of wild and laboratory strains of *Rattus norvegicus* (F-344, Wistar) eight weeks after inoculation with various sporocyst doses (400–10,000). Bradyzoites were obtained by tryptic digestion of muscles; values are the mean \pm standard deviation of 6–8 rats per treatment.

In conclusion, *S. singaporensis* infection in rodents induces almost 100% mortality once above a threshold level of infection (Table 3). The effect of *S. singaporensis* infection on the fertility of rodents is less well founded. When there is a reduction in fertility, the effect may be influenced by the sex and age

of the host, its genetic background, or status of the immune system.

***S. singaporensis* as a potential biological control agent**

We have examined whether this parasite could be an effective tactical tool for rodent control by artificially disseminating food pellets containing sporocysts among rodents in the field.

Evidence that the parasite increases the mortality of rodents under natural conditions was provided during a field experiment performed in Egypt (Jäkel et al. 1996). Infection of a small population of roof rats (*Rattus rattus frugivorous*) with food-pellets containing a lethal amount of sporocysts killed 73% of the rats when

compared to the effect of a placebo. A characteristic time-course of activity of rodents artificially infected in the field is presented in Figure 5.

Recent field experiments in Thailand (plots up to 4 ha) showed that *S. singaporensis* is highly effective against *R. norvegicus* and *B. indica*. Parasite-induced mortality ranged between 60% and 80% (Jäkel et al. 1997a; Jäkel et al., data submitted for publication). Importantly, the latter results indicate that *S. singaporensis* can be used as a biocontrol agent inside its natural distribution range in Southeast Asia despite the fact that the parasite frequently occurs in rodents in this region (O'Donoghue et al. 1987; Jäkel et al. 1997b). This conforms with the prospects previously outlined by Wood (1985).




Figure 5. Representative measurement of the activity of wild rats (*Rattus norvegicus*) after infection with *Sarcocystis singaporensis* (day 0 = day of infection) (modified from Jäkel et al. 1996, 1997a). The activity of rats (10–100 animals) is expressed as the consumption of plain bait or the number of footprints on tracking plates. Note that activity declines around day 10, indicating the onset of parasite-induced mortality among rats. Usually, dead rats can be seen in the field 10–16 days post infection. At that time, merozoites of *S. singaporensis* synchronously leave their host cells in the lungs inducing a fatal pneumonia in their hosts.

Laboratory experiments showed that natural infections usually do not provide protection against lethal sporocyst doses (Table 3). The reasons behind this remain to be fully determined, however, analysis of the T-cell response after infection revealed that low numbers of sporocysts do not induce the formation of memory CD4 T-cells (Jäkel et al. 1998) which are usually responsible for the persistence of immuno-logical memory in the rat (Bell et al. 1998). Assuming that the high numbers of bradyzoites developing in the muscles (Figure 4) are proportional to those of the preceding stage (the merozoite which can cause extensive pathology), these data indicate that rats can tolerate substantial numbers of multiplying parasites before they show signs of disease. A certain degree of infection is tolerated, and may even suppress immune function (Gill et al. 1988). However, rats can become resistant to acute infection if they ingest a sublethal but high number of sporocysts (Jäkel et al. 1996); accordingly, in this case numbers of memory CD4 T-cells were significantly increased (Jäkel et al. 1998). With regard to rodent control, these observations indicate that low numbers of parasites in the environment are no obstacles to control measures using the parasite. However, a proper bait formulation is needed which prevents unintended immunisations during field application.

From research into practice

Current research activities of the project focus on three major topics: (i) development of a parasite-bait which is highly palatable and preserves viability of sporocysts, (ii) developing conditions for mass-production of the parasite, and (iii) defining an application scheme which could complement

or become integrated with other practices of rodent control.

Developing a suitable parasite-bait is the most difficult part and will be the key factor in a possible commercial exploitation of the method. During our field experiments, a bait was used which had been developed by the Bayer AG, Monheim, Germany. It consisted of a wheat paste with a high oil content and was sweetened with sugar. The parasite-inoculated mixture was superior to other food items and especially attractive to *Rattus* spp. and *B. indica* from Thailand. Unfortunately, a high oil content seems to have negative effects on the viability of sporocysts, and therefore other parasite-bait formulations are currently under investigation. The project aims to achieve a storage stability period of at least three months under local conditions (high ambient temperature, high humidity).

Sporocysts can be mass-produced in reticulated pythons (Jäkel et al. 1996), therefore the project is constructing a pilot production unit in cooperation with Thai private industry to rear and keep pythons. In Thailand, snake farms are found all over the country, serving as tourist attractions or providing products for the leather industry. We have observed experimentally that pythons (2–3 m length) shed up to 4×10^9 sporocysts in the faeces after a single infection without showing any signs of disease. The animals can be reinfected six to eight times per year. In the snake's intestine, immune reactions against the parasite are not apparent. A lethal dose for a single rat ranges between 2×10^5 sporocysts (inside the natural distribution range of the parasite; Table 3) to 2×10^4 sporocysts (outside the natural distribution range; Jäkel et al. 1996).

Therefore a single infection of a snake can yield material to kill about 2×10^4 to 2×10^5 rats.

S. singaporensis could be a new tactical tool in rodent control, with its application being similar to chemical rodenticides, and complementing other non-chemical approaches (McCallum 1996; Chambers et al. 1997; Singleton et al. 1998). Field studies are planned to determine the effectiveness and acceptance of the method at the farmers' level. As well as an easy-to-use design of a parasite-bait, a low price will be crucial for its success on the rodenticide market.

SYNOPSIS AND FUTURE CONCEPTS

As outlined previously, rodent management in Thailand mainly relies on the use of chemical rodenticides and mechanical methods, an approach which was developed in the mid-seventies. This approach has also contributed to a substantial reduction in the rodent problem, especially in rice, because extension programs were continuously conducted to reach the farmer. However, rodent damage to agriculture continues, notably that by mice to various field crops or climbing rat species to oil palm. Effective control against rats is lacking in crop stores of small landholders or in urban areas. Recent outbreaks of leptospirosis among humans in Thailand indicate that research and practical control measures are necessary to restrict the spreading of rodent-borne diseases.

According to the 8th National Economic and Social Development Plan (1997–2001), it is the policy of the Ministry of Agriculture and Cooperatives that the use of pesticides should be reduced as much as possible.

Therefore, efforts are underway to implement an integrated pest management (IPM) strategy in selected pilot areas in the country. In 1997, the Department of Agricultural Extension started an IPM program against plant diseases and insect pests including 100 demonstration plots (80 ha each) in various parts of the country. Currently, strategies for rodent control management included in the curricula of the education program mainly focus on chemical approaches, with environmentally friendly techniques only playing a minor role at present. The AZRG is testing new methods in biological control using the parasitic protozoan *Sarcocystis singaporensis* and mechanical approaches like the trap-barrier system. It is planned to regularly apply these techniques in demonstration plots of the above mentioned IPM program. The future will show if these methods are accepted by farmers in Thailand and can be promoted at a larger scale.

REFERENCES

- Anonymous 1989. Evaluation of adoption by farmers of rodent control practices in the Rodent Control Campaign Project. Division of Project and Program Evaluation Office of Agricultural Economics. Ministry of Agriculture and Cooperatives, 59p. (in Thai).
- Anonymous 1996. Agricultural data at provincial level. Division of Planning, Department of Agricultural Extension, 160p. (in Thai).
- Artchawakom, T., Hongnark, S., Khoprasert, Y. and Chanyapate, C. 1986. Study on rodent pests and damage appraisal of barley. Research Annual Report, Entomology and Zoology Division, Department of Agriculture, 1–4 (in Thai).
- Beaver, P.C. and Maleckar, J.R. 1981. *Sarcocystis singaporensis*, *Sarcocystis villivillosi* sp. n., and *Sarcocystis zamani* sp. n.: development,

- morphology and persistence in the laboratory rat, *Rattus norvegicus*. The Journal of Parasitology, 67, 241–256.
- Bell, E.B., Sparshott, S.M. and Bunce, C. 1998. CD4+ T-cell memory, CD45R subsets and the persistence of antigen—a unifying concept. Immunology Today, 19, 60–64.
- Boonsong, P., Chanyapate, C., Artchawakom, T. and Somsook, S. 1984a. Observation of reproductivity of rice pest rats in Amphor Banglan Nakornpathom. Research Annual Report, Entomology and Zoology Division, Department of Agriculture, 19p. (in Thai).
- Boonsong, P., Chaowattanawong, P., Dhammabamrung, N., Hongnark, S., Artchawakom, T., Sianglew, P. and Sarakun, N. 1987. Assessment of damage caused by rats in oil palm. Research Annual Report. Entomology and Zoology Division, Department of Agriculture, 11p. (in Thai).
- Boonsong, P., Hongnark, S., Niyomvit, L. and Khoprasert, Y. 1994. Survey on rodenticides used by farmers in soybean fields in 1994. Research Annual Report. Entomology and Zoology Division, Department of Agriculture, 136–158 (in Thai).
- Boongsong, P., Puangprakhone, K., Khoprasert, Y., Artchawakom, T., Chanyapate, C. and Chamkrachang, W. 1991. Population dynamic study of rodents in maize. Research Annual Report, Entomology and Zoology Division, Department of Agriculture (in Thai), 154–167.
- Boonsong, P., Somsook, S., Artchawakom, T., Seehabutr, V., Suwanachai, C. and Tongtavee, K. 1984b. Rat movement in rice fields. Research Annual Report, Entomology and Zoology Division, Department of Agriculture, 1–10 (in Thai).
- Boonsong, T., Niyomvit, L., Hongnark, S., Khoprasert, Y. and Nookarn, P. 1995. Survey on rodenticides used by farmers in soybean fields in the lower north of Thailand. Research Annual Report. Entomology and Zoology Division, Department of Agriculture, 91–133 (in Thai).
- Boonsong, T., Niyomvit, L., Hongnark, S., Nookarn, P. and Chanyapate, C. 1996. Survey on rodenticides used by farmers in soybean fields in the northeast of Thailand. Abstract in Research Annual Report, Entomology and Zoology Division, Department of Agriculture, 47–50 (in Thai).
- Brehm, H. and Frank, W. 1980. Der Entwicklungskreislauf von *Sarcocystis singaporensis* Zaman und Colley, 1976 im End- und Zwischenwirt. Zeitschrift für Parasitenkunde, 62, 15–30.
- Chambers, L.K., Singleton, G.R. and Hood, G.M. 1997. Immunocontraception as a potential control method of wild rodent populations. Belgian Journal of Zoology, 127 (suppl. 1), 145–156.
- Chokvivat, W. 1998. Leptospirosis. Department of Communicable Disease Control, Ministry of Health. Bangkok, Cooperative Press, 104p.
- Corbet, G.B. and Hill, J.E. 1992. The mammals of Indomalayan region: a systematic review. Natural History Museum Publications. Oxford, Oxford University Press, 448p.
- Fayer, R., Andrews, C. and Dubey, J.P. 1988. Lysates of *Sarcocystis cruzi* bradyzoites stimulate RAW 264.7 macrophages to produce tumor necrosis factor (cachectin). The Journal of Parasitology, 74, 660–664.
- Gill, H.S., Charleston, W.A.G. and Moriarty, K.M. 1988. Immunosuppression in *Sarcocystis muris*-infected mice: evidence for suppression of antibody and cell-mediated responses to a heterologous antigen. Immunology and Cell Biology, 66, 209–214.
- Gratz, N.G. 1994. Rodents as carriers of disease. In: Buckle, A.P. and Smith, R.H., ed., Rodent pests and their control. Wallingford, UK, CAB International, 85–108.
- Häfner, U. 1987. Zystenbildende Coccidien mit Nager/Schlange-Syklen unter besonderer Berücksichtigung der Wirtsspezifität der Gattung *Sarcocystis*. Dissertation, Universität Hohenheim, Stuttgart.
- Häfner, U. and Frank, W. 1984. Host specificity and host range of the genus *Sarcocystis* in three snake-rodent life cycles. Zentralblatt für Bakteriologie, Mikrobiologie und Hygiene, Originale A, 256, 296–299.
- Hongnark, S., Boonsong, P., Khoprasert, Y., Tippayarak, S., Tangjitrong, A. and Jairin, J. 1994. Study on rodent pests and damage

- appraisal of wheat. Research Annual Report, Entomology and Zoology Division, Department of Agriculture, 75–81 (in Thai).
- Hongnark, S., Boonsong, T., Tongtavee, K., Chanyapate, C., Suasa-ard, K. and Kaewta, T. 1993. Damage appraisal of rice due to rodent pests in Thailand. Research Annual Report, Entomology and Zoology Division, Department of Agriculture, 10–18 (in Thai).
- Hongnark, S., Sudto, P., Wongraj, C., Artchawakom, T., Somsook, S. and Tongtavee, K. 1984. Study on bird pests, damage appraisal and their control in upland rice. Research Annual Report, Entomology and Zoology Division, Department of Agriculture, 9p. (in Thai).
- Hoogenboom, I. and Dijkstra, C. 1987. *Sarcocystis cernae*: a parasite increasing the risk of predation of its intermediate host, *Microtus arvalis*. *Oecologia*, 74, 86–92.
- Jäkel, T., Burgstaller, H. and Frank, W. 1996. *Sarcocystis singaporensis*: studies on host specificity, pathogenicity, and potential use as a biocontrol agent of wild rats. *The Journal of Parasitology*, 82, 280–287.
- Jäkel, T., Khoprasert, Y., Baumann, C., Kliemt, D., Sangchai, J. and Hongnark, S. 1997a. Artificial *Sarcocystis* infection in controlling commensal rodents. Abstract, 7th International Theriological Congress, September 6–11, Acapulco, Mexico.
- Jäkel, T., Khoprasert, Y., Sorger, I., Kliemt, D., Seehabutr, V., Suasa-ard, K. and Hongnark, S. 1997b. *Sarcosporidiosis* in rodents from Thailand. *Journal of Wildlife Diseases*, 33, 860–867.
- Jäkel, T., Sangchai, J. and Khoprasert, Y. 1998. The role of immunological memory in *Sarcocystis* infection. In: Ortega-Mora, L.M., Gomez-Bautista, M., Bueno-Pereira, J. and Tenter, A.M., ed., Proceedings of the COST-820 1998 Annual Workshop: vaccines against animal coccidiosis. Toledo, Spain, European Commission, 50.
- Khoprasert, Y., Artchawakom, T., Sehabutr, V. and Boonsong, P. 1990. Survey on rat species and damage assessment in longan. Research Annual Report, Entomology and Zoology Division, Department of Agriculture, 62–68 (in Thai).
- Khoprasert, Y., Suasa-ard, K., Chanyapate, C., Wongraj, C., Suwanachai, C. and Thitipawat, K. 1977. Movement study of rodents in the rice fields. Research Annual Report, Entomology and Zoology Division, Department of Agriculture, 2p. (in Thai).
- Lietmeyer, K. 1988. Epizootiology of hantaviruses in northern Thailand. Masters Thesis of Public Health, Yale University.
- McCallum, H. 1996. Immuncontraception for wildlife population control. *Trends in Ecology and Evolution*, 11, 491–493.
- O'Donoghue, J.P., Watts, C.H.S. and Dixon, B.R. 1987. Ultrastructure of *Sarcocystis* spp. (*Protozoa apicomplexa*) in rodents from North Sulawesi and West Java, Indonesia. *Journal of Wildlife Diseases*, 23, 225–232.
- Prakash, I. 1988. Bait shyness and poison aversion. In: Prakash, I., ed., *Rodent pest management*. Boca Raton, CRC Press, 321–329.
- Ratanaworabhan, S. 1971. Rodent pests and their control. *Journal of Agricultural Science*, 4, 45–57 (in Thai).
- Rommel, M. and Heydorn, A.O. 1972. Beitrage zum Lebenszyklus der Sarcosporidien, III. *Isospora hominis* (Railliet und Lucet, 1891) Wenyon, 1923, eine Dauerform der Sarcosporidien des Rindes und des Schweinsl. *Berliner und Müncher Tierärztliche Wochenschrift*, 85, 143–145.
- Rzepczyk, C. and Scholtyseck, E. 1976. Light and electron microscope studies on the *Sarcocystis* of *Rattus fuscipes*, an Australian rat. *Zeitschrift für Parasitenkunde*, 50, 137–150.
- Schmaljohn, C. and Hjelle, B. 1997. Hantaviruses: a global disease problem. *Emerging Infectious Diseases*, 3, 95–104.
- Schrag, S.J. and Wiener, P. 1995. Emerging infectious disease: what are the relative roles of ecology and evolution? *Trends in Ecology and Evolution*, 10, 319–324.

- Silapapochakul, K. 1992. Acute PUO. In: Infectious Disease Society of Thailand: current therapy of common infectious disease. Bangkok, Thailand, Medical Media, 66–78.
- Singleton, G.R. and Petch, D.A. 1994. A review of the biology and management of rodent pests in Southeast Asia. Canberra, Australian Centre for Agricultural Research, Technical Report No. 30, 65p.
- Singleton, G.R., Sudarmaji and Suriapermana, S. 1998. An experimental field study to evaluate a trap–barrier system and fumigation for controlling the rice field rat, *Rattus argentiventer*, in rice crops in West Java. *Crop Protection*, 17, 55–64.
- Somsook, S., Hongnark, S., Suwanachai, C., Kaewta, T., Artchawakom, T. and Tongtavee, K. 1983. Study on population dynamics of *Rattus losea* (Swinhoe) in rice fields. Research Annual Report. Entomology and Zoology Division, Department of Agriculture, 10p. (in Thai).
- Suasa-ard, K., Hongnark, S., Khoprasert, Y., Sawanachai, S., Bamrungsuk, S. and Keawta, T. 1987. Survey on rat species and damage assessment in sugar cane fields. Research Annual Report, Entomology and Zoology Division, Department of Agriculture, 20p. (in Thai).
- Weis, N. 1981. Rodent pests and their control. Eschborn, Germany, GTZ (Deutsche Gesellschaft für Technische Zusammenarbeit), 206p.
- Wood, B.J. 1985. Biological control of vertebrates—a review, and assessment of prospects for Malaysia. *Journal of Plant Protection in the Tropics*, 2, 67–97.
- Wood, B.J. 1987. Oil palm pest control in Thailand. Thailand Oil Palm Research and Development Project THA/84/007, FAO. Kuala Lumpur, Sime Darby Services, 82p.
- Wood, B.J. 1994. Rodents in agriculture and forestry. In: Buckle, A.P. and Smith, R.H., ed., *Rodent pests and their control*. Wallingford, UK, CAB International, 45–83.
- Wood, B.J. and Liao, S.S. 1984. A long term study of *R. tiomanicus* populations in an oil palm plantation in Johore, Malaysia. II. Recovery from control, and economic aspects. *Journal of Applied Ecology*, 21, 465–472.
- Zaman, V. 1976. Host range of *Sarcocystis orientalis*. *Southeast Asian Journal of Tropical Medicine and Public Health*, 7, 112.
- Zaman, V. and Colley, F.C. 1975. Light and electron microscopic observations of the life cycle of *Sarcocystis orientalis* sp. n. in the rat (*Rattus norvegicus*) and the Malaysian reticulated python (*Python reticulatus*). *Zeitschrift für Parasitenkunde*, 47, 169–185.